



Comparative single-cell analysis of transcriptional bursting reveals the role of genome organization in de novo transcript origination

Unjin Lee^a , Cong Li^a , Christopher B. Langer^a, Nicolas Svetec^a , and Li Zhao^{a,1}

Edited by Harmit Malik, Fred Hutchinson Cancer Center, Seattle, WA; received December 11, 2024; accepted April 2, 2025

Spermatogenesis is a key developmental process underlying the origination of newly evolved genes. However, rapid cell type-specific transcriptomic divergence of the *Drosophila* germline has posed a significant technical barrier for comparative single-cell RNA-sequencing studies. By quantifying a surprisingly strong correlation between species- and cell type-specific divergence in three closely related *Drosophila* species, we apply a statistical procedure to identify a core set of 198 genes that are highly predictive of cell type identity while remaining robust to species-specific differences that span over 25 to 30 My of evolution. We then utilize cell type classifications based on the 198-gene set to show how transcriptional divergence in cell type increases throughout spermatogenic developmental time. After validating these cross-species cell type classifications using RNA fluorescence in situ hybridization and imaging, we then investigate the influence of genome organization on the molecular evolution of spermatogenesis vis-a-vis transcriptional bursting. We first show altering transcriptional burst size contributes to premeiotic transcription and altering bursting frequency contributes to postmeiotic expression. We then report global differences in autosomal vs. X chromosomal transcription may arise in a developmental stage preceding full testis organogenesis by showing evolutionarily conserved decreases in X-linked transcription bursting kinetics in all examined somatic and germline cell types. Finally, we provide evidence supporting the cultivator model of de novo gene origination by demonstrating how the appearance of newly evolved testis-specific transcripts potentially provides short-range regulation of neighboring genes' transcriptional bursting properties during key stages of spermatogenesis.

single-cell RNA-seq | transcriptional divergence | de novo genes | *Drosophila* | testis

Gametogenesis is an essential process in all sexually reproducing species in nature, ensuring genetic material is successfully transferred between generations. Male gametogenesis, spermatogenesis, is a specialized developmental process that exhibits several key features. First, cell differentiation from stem cell to fully formed gamete is remarkably conserved across animals. Second, despite the necessary maintenance of gametic function, testis-expressed genes tend to evolve very rapidly, with some genes being newly originated. Third, although cell types in the testis exhibit conserved features, expression patterns of genes in each cell type vary drastically between species. Because of this, the expression dynamics of testis-expressed genes are an important issue for both evolutionary biology and developmental biology.

While the transcriptomes of highly conserved cell types in the testis are likely to remain stable over large evolutionary timescales, the rapid evolution of spermatogenesis in *Drosophila* presents significant challenges for comparative analyses of single-cell transcriptomic data, even among closely related species. For instance, many seemingly conserved marker genes from *Drosophila melanogaster* (1, 2) are lost or not annotated in other *Drosophila* species, challenging the assumption that these genes are evolutionarily conserved. Additionally, the rate of gene expression alteration drastically varies in *Drosophila* (3, 4), making it difficult to transfer cell types across different species. Our interest lies not only in the conserved transcriptional programs of cell types across species but also in complete gene expression patterns, including species-specific genes. The application of standard techniques (5–8) often fails to reproduce previous cell-type assignments in *D. melanogaster* samples, generating a large number of species-specific cell-type clusters that likely do not reflect true phenotypic differences.

Within approximately 30 My of evolution in the *melanogaster* species group, the process of germline differentiation has undergone many phenotypic changes (9–11). Alternatively, a large degree of spermatogenesis (Fig. 1*A*) has remained unaltered through similar evolutionary time scales (12, 13), including gross anatomical phenotypes such as organ structure or the developmental progression of various cell-type specific changes (14, 15). We

Significance

Understanding the divergence and evolution of novel genes and expression is an essential question in evolutionary biology. However, rapid transcriptomic divergence in tissues such as the testis has significantly hindered comparative single-cell RNA-Sequencing (scRNA-Seq) studies. Here, we provide a strategy— independent of preexisting cell type marker genes—to overcome this barrier. In finding how genome organization influences the evolution of transcriptional bursting dynamics via de novo transcript origination, our results support the cultivator model of de novo gene origination—a model emphasizing the genomic environment in shaping novel gene origination.

Author affiliations: ^aLaboratory of Evolutionary Genetics and Genomics, The Rockefeller University, New York, NY 10065

Author contributions: U.L. and L.Z. designed research; U.L., C.L., C.B.L., N.S., and L.Z. performed research; U.L., C.L., C.B.L., and N.S. contributed new reagents/analytic tools; U.L. analyzed data; and U.L. and L.Z. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Copyright © 2025 the Author(s). Published by PNAS. This article is distributed under Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND).

¹To whom correspondence may be addressed. Email: lzhao@rockefeller.edu.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2425618122/-DCSupplemental>.

Published April 30, 2025.




Fig. 1. Identification of 198-gene list allows for cell type classification across species in a rapidly diverging process. (A) Illustration of spermatogenesis from germline stem cells to late spermatids. (B) Phylogeny of species used in this study with estimated divergence times. (C) ANOVA $\log(F\text{-statistic})$ values from scRNA-Seq data using previously identified *D. melanogaster* cell types and cross-species labels for the three species show high correlation resulting in a large degree of species-specific cell type divergence. Drawn lines correspond to median $F\text{-statistic}$ values for each (independent) marginal distribution. (D–F) Uniform Manifold Approximation and Projections (UMAPs) resulting from the 198-gene list reveal a progression of intermediate cell types spanning germline stem cells to late spermatids for the three species. (G) Violin plots from the most differentially expressed gene within each cell type.

therefore hypothesized that a high degree of species-specific gene expression evolution within each cell type is the primary factor driving technical difficulties in clustering and identifying cell type differences within comparative studies of *Drosophila* spermatogenesis (SI Appendix, Fig. S1).

Transcription is an intrinsically random process that may be modeled stochastically (16, 17). Single-molecule observations of transcriptional activity are intrinsically “bursty,” consisting of short bouts of transcription that produce discrete quantities of mRNAs (“on” state) followed by periods of inactivity (“off” state) (18–20). Two key parameters controlling the transcriptional kinetics of different genes are burst size (or burst amplitude), which reflects the number of transcripts produced per burst, and burst frequency, which reflects the number of bursts that occur per unit time (18, 21, 22). Importantly, the analysis of transcriptional bursting kinetics offers unique insight into understanding how certain genes are regulated. The kinetics underlying burst size is a property of core

promoter elements, driven by the stoichiometry of RNA polymerase II (pol II) availability, while the kinetics of burst frequency is influenced primarily by enhancer elements, modulated by the probability of transcriptional initiation (23–26). Transcriptional bursting properties have provided a novel perspective into the molecular underpinnings of sex-biased transcription burst amplitude in male and female embryos of *D. melanogaster* (27). Recent work has shown that the upregulation of male X chromosome genes is primarily mediated by a higher pol II initiation rate and burst amplitude (27), paving the way for further studies of dosage compensation in spermatogenesis.

Despite this, the molecular mechanisms underlying downregulation of the X chromosome remain understudied (28). While prior work has suggested that expression of genes on the X chromosome is globally down-regulated during spermatogenesis in *Drosophila* (29), it is not possible to determine anything beyond the general reduction of expression for X-linked genes (28, 30).

Alternatively, previous single-cell RNA-sequencing (scRNA-Seq) studies have proposed that there is some degree of noncanonical dosage compensation (31, 32). A decrease in X chromosome expression beyond that which would be expected by dosage has also led others to suggest that the entire *Drosophila* X chromosome is inactivated (33).

Beyond the X chromosome, our interest in the testis is also related to the observation that the testis is a hotspot for tissue-specific expression of newly evolved genes (34, 35), including novel genes and transcripts (e.g., noncoding RNAs) arising de novo from nongenic, transcriptionally inactive sequences. While the origination of de novo protein-coding genes has been relatively well-studied (36–38), the study of the evolutionary mechanisms underlying the origination of de novo transcripts, e.g., long noncoding RNA (lncRNA), in fruit flies has been lacking (36, 39, 40). This gap highlights the importance of comprehensively identifying these RNA genes in multiple *Drosophila* species (41). Such de novo transcripts are often expressed in a tissue-specific manner, typically in the adult testis of *Drosophila* (34, 36, 39). Intriguingly, scRNA-Seq data have shown that de novo originated genes, unlike recently evolved duplicate genes, are biased toward expression in mid-spermatogenesis (35). It is not yet clear whether this pattern is common across multiple *Drosophila* species and what function these transcripts may hold.

In our study, we undertook a comprehensive analysis of spermatogenesis in three *Drosophila* species, aiming to unravel the evolution of complex transcriptional dynamics and novel gene expression. We first developed an approach to study scRNA-Seq data from rapidly diverging species. We then quantified a strong correlation between species- and cell-type-specific transcriptomic divergence across these species, identifying a core set of genes crucial for predicting cell type identity over approximately 30 My of evolutionary divergence. After classifying cell types across these species, we analyzed how transcriptional divergence progresses throughout spermatogenic development. We further investigated the effects of genome organization on the molecular evolution of spermatogenesis, focusing on transcriptional bursting. We demonstrated variations in transcriptional burst size and frequency before and after meiosis, examining differences in autosomal versus X chromosomal transcription. Our findings support the cultivator model of de novo gene origination (42), showing that newly evolved testis-specific transcripts might play a regulatory role in the transcriptional bursting properties of neighboring genes.

Results

A Simple ANOVA-Based Statistical Procedure Identifies Gene Lists Predictive of Cell Type but Not Species Identity. We used several commonly available software suites to analyze scRNA-Seq data from testis tissue derived from three species, *D. melanogaster*, *Drosophila yakuba*, and *Drosophila ananassae*, but were unable to obtain high-quality cell type assignments to correlate similar cell types across different species (SI Appendix, Fig. S1). For example, widely used markers in *D. melanogaster* did not robustly classify cell type identities in species such as *D. ananassae*. We thus hypothesized that species-specific cell type transcriptomic divergence significantly hinders cell type classification. Clear morphological and functional conservation of spermatogenesis across various *Drosophila* species, e.g., mitotic staging or spermatid differentiation, suggests some degree of evolutionary conservation of key genetic elements. In this case, challenges in cross-species clustering could potentially be overcome by focusing on key conserved elements, not the entire transcriptome. We thus developed a simple methodology to identify genes that are both evolutionarily conserved and important in determining cell identity. We hypothesized that these genes could

be identified by calculating ANOVA F -statistic scores for cell type (where available) and species. To test this strategy, we generated a simulated dataset consisting of nine unique cell types evolving in three different species (SI Appendix, Fig. S2A), to simulate a scenario in which cell type differences are masked by species-level differences. (SI Appendix, Fig. S2A). A good cell type marker gene, indexed by variable i , for multiple species should exhibit a large F_i for cell type and a small F_i for species identity (SI Appendix, Fig. S2B, Bottom-Right quadrant). We selected the top 200 genes showing high F_i (cell type) and low F_i (species identity). The genes identified through our methodology effectively captured the most informative simulated genes, preferentially containing cell type information over species identity information (SI Appendix, Fig. S2C).

Uncorrected simulated data resulted in 27 nonoverlapping cell types, as they were clustered based on both cell type and species identity (SI Appendix, Fig. S2D). When standard batch correction techniques, such as Monocle3's implementation of batchelor via "align_cds()" (6, 43), are applied to remove species-specific effects, Uniform Manifold Approximation and Projection (UMAP) and Principle Component Analysis (PCA) projections show that these effects have been removed with entirely overlapping cell type assignments despite species-specific differences in cell type (SI Appendix, Fig. S2E). This stands in contrast to the removal of species-specific changes using these same batch correction techniques in real scRNA-Seq data derived from testis tissue (SI Appendix, Fig. S1), as remaining species-specific differences hinder the identification of cell types across species in these datasets. This is likely due to the excessively simplistic and additive nature of the simulations in contrast to the more complex, nonlinear nature of cell-type specific evolution. Regardless, the application of standard batch correction techniques remained insufficient for cell type identification in real data.

Projection and clustering using the top 200 gene list in the simulated data showed nine distinct cell type clusters that still maintained species-specific differences (SI Appendix, Fig. S2F and G). Cell type identities are readily recovered, while species-specific divergence within these cell types is also clearly visualized. These show that our method identifies gene lists that are predictive of cell type but not species identity, which is important for clustering cell types across species. The methodology is also effective across phylogenies of varying degrees of asymmetry (SI Appendix, Fig. S3 and Performance of H_{SQ} Criteria in Asymmetric Phylogenies).

Application of the ANOVA-Based Statistical Procedure Identifies 198 Gene List in Related *Drosophila* Species. We then applied our nonparametric statistical procedure on our scRNA-Seq dataset derived from testis tissue in *D. melanogaster*, *D. yakuba*, and *D. ananassae*. The divergence time between *D. melanogaster* and *D. yakuba* has been estimated to be about 7 Mya while the divergence time between *D. melanogaster* and *D. ananassae* has been estimated to be about 25 to 30 Mya (44) (Fig. 1B). Prior data and cell type labels from *D. melanogaster* were obtained from ref. 35, while 5,000 cells derived from *D. yakuba* and 5,000 cells derived from *D. ananassae* were sequenced (SI Appendix). Raw data were aligned in "cellranger" to reference genomes and annotations obtained from FlyBase to generate read counts. Further orthology information between the three species' annotations was obtained using FlyBase (dmel-r6.44) and used to generate "melanogasterized" *D. yakuba* (dyak-r1.05) and *D. ananassae* (dana-r1.06) genomes. Specifically, each genome was reduced to a core set of single-copy genes that possessed strict one-to-one orthology between the three species. After processing in "Monocle3" (43), data from one-to-one orthologs were combined together and retained. A summary of key processing parameters is presented in SI Appendix, Table S1.

The distributions of F_i (cell type) and F_i (species identity) were calculated using our experimentally derived data (Fig. 1C and Dataset S1). Interestingly, the joint distribution of $\log(F_i$ (cell type)) and $\log(F_i$ (species identity)) showed an exceedingly large correlation (Pearson $\rho = 0.698$, $P < 2 \times 10^{-16}$, Fig. 1C). This result is on par with the correlation found in simulated data (SI Appendix, Fig. S2B, Pearson $\rho = 0.74$, $P < 2 \times 10^{-16}$), suggesting a high degree of species-specific cell type evolution in *Drosophila* testis tissue. The medians of the independent distributions F_i (cell type) and F_i (species identity) were chosen as threshold $p_{\text{cell-type}}$ and $p_{\text{species-identity}}$ respectively (Fig. 1C, solid lines). Note that the independent distributions of F_i (cell type) and F_i (species identity) are not fully represented in the joint distribution of F_i (cell type) and F_i (species identity), because some genes were missing F_i (cell type) or F_i (species identity) values. This definition of $p_{\text{cell-type}}$ and $p_{\text{species-identity}}$ produced a new 198-gene (SI Appendix, Table S2).

To support the biological significance of the 198-gene set, we performed a gene ontology analysis using the PANTHER database (45, 46) (SI Appendix, Table S3). The 11 statistically significant major categories (FDR < 0.05) were highly concordant with known function during spermatogenesis, including “cilium mobility,” “dynein-associated pathways,” “insulin signaling,” “centrosome cycle,” “protein maturation,” “proteolysis,” and “general reproduction.” Notably, all of these major categories showed overrepresentation in comparison to what is expected for all genes in *D. melanogaster*, demonstrating that this statistical procedure effectively identifies transcriptionally important, functionally conserved genes in the rapidly evolving *Drosophila* testis transcriptome.

Cell Type Clustering of Testis scRNA-Seq Data Reveals Increasing Transcriptomic Divergence Throughout Spermatogenesis. To validate the expression of the 198-gene set we identified, we visualized both UMAP and PCA projections of all analyzed cells with an overlay of previously identified cell type assignments (35) (SI Appendix, Fig. S4A). Importantly, cells from all species can be projected onto the same visualization (subsetting by the 198-gene list) without extraneous processing beyond standard single-cell transcriptomics pipelines (Fig. 1D–F and SI Appendix, Fig. S4B). We note that batch correction for species was applied to the simulated 200- and this 198-gene list as part of Monocle3’s standard pipeline. A visual examination of these projections shows striking features concordant with known spermatogenesis function. 1) While the sequence of spermatogenesis was not explicitly encoded into our statistical procedure or in the visualization process, the developmental progression of cell types in the projection follows a known developmental progression from germline stem cells to late spermatids. Despite being a continuous developmental process where each stage does not produce spatially segregated clusters, the stages of previous cell type assignments form relatively distinct boundaries. This eliminates the need for more advanced pseudotime projection techniques. 2) All somatic cell types cluster separately from germ cell types. 3) We observe a gradual increase in cell-to-cell gene expression variance as development progresses. For example, germline stem cells appear to have low expression variance across all species, while spermatids show much larger variance (Fig. 1D–F). Note that all these features are readily observed in PCA projections of the same dataset (SI Appendix, Fig. S4B).

Importantly, the application of this methodology allows for recapitulation of previously identified cell type clusters using unsupervised clustering on a unified, panspecies UMAP with the clustering resolution set to produce the same number of clusters reported in previous studies (35) (Leiden clustering, resolution = 3×10^{-4} , nine clusters). While it is likely to be the result of residual

species-specific differences, these cell type assignments show that fewer cells are represented in the earlier germline stem cell, spermatogonia, and spermatocyte stages of *D. yakuba* when compared to *D. melanogaster*. We also observe a correlated accumulation of cells in later spermatid developmental stages. These cell type assignments also reveal the presence of an additional cell cluster/ expression present in *D. ananassae* that appears to be absent from *D. melanogaster* and *D. yakuba*. We believe that this cluster is likely a result of transcriptomic divergence (over evolutionary timescales) that persists even after batch correction. Specifically, these cells likely represent a set of spermatocyte cells that demonstrate a stereotypical expression pattern that exists only in *D. ananassae* (New Spermatocyte Cell Cluster Reveals a Unique Expression Pattern and SI Appendix: ‘ananassae’ spermatocyte cluster). We also observe a low abundance of early and late spermatid stages in *D. ananassae* (Fig. 1D–F and SI Appendix, Fig. S4B). These changes in cell type assignment and visualized patterns among species reflect changes in the expression patterns of the 198-gene list, possible technical biases during data generation, and fundamental processes driving species-specific differences in reproduction (New Spermatocyte Cell Cluster Reveals a Unique Expression Pattern and SI Appendix: ‘ananassae’ spermatocyte cluster). These cell clusters were further investigated using RNA fluorescence in situ hybridization (FISH) (RNA In Situ Hybridization Chain Reaction Validates Cross-species Cell Type Assignments.)

High Differential Expression across Cell Types in *D. melanogaster* Spermatogenesis Does Not Necessarily Indicate Differential Expression across Cell Types in Closely Related Species. A cross-species analysis of the five most significantly differentially expressed genes for each cell type/cluster was performed. Unless specifically noted, the phrase “differential expression” indicates analyses of differential expression across cell types that do not take species identity into account. For example, if a gene is said to be differentially expressed in Germline stem cell (GSC)/Early spermatogonia, we mean that the expression of this gene is altered when comparing GSC/Early spermatogonia for all species to all other cell types for all species. This is achieved by combining GSC/Early spermatogonia from *D. melanogaster*, *D. yakuba*, and *D. ananassae* into one category and comparing these cells to all other cells detected in *D. melanogaster*, *D. yakuba*, and *D. ananassae*.

Interestingly, only one (*MtnA*) of 13 previously reported markers (*aly*, *zfh1*, *Fas1*, *Hsp23*, *MtnA*, *His2Av*, *aub*, *bam*, *fzo*, *twe*, *soti*, *Dpy-30L2*, *p-cup*) (35) remains as one of the top-five differentially expressed genes by cell type (all spp., e.g. all GSC from all spp. considered together, SI Appendix, Fig. S5). These results suggest that while the 13 marker genes identified in previous *D. melanogaster* studies may be highly differentially expressed across cell types in *D. melanogaster*, these expression patterns are likely to have evolved in a species-biased manner. A lack of one-to-one orthology may also be excluded as a driving factor in these discrepancies, as only one of these 13 markers (*zfh1*) was removed due to this reason.

We then analyzed all significantly differentially expressed genes for each cell type, testing for both over- and underexpression (SI Appendix, Table S4 and Dataset S2). This analysis was performed by comparing cell types without regard to species. Among the 12 previously used markers present in the dataset, *Fas1*, *fzo*, and *Dpy-30L2* were not found to be differentially expressed in any of these cell types (all spp.). In the somatic cells, *Hsp23* and *MtnA* remained highly differentially expressed (all spp.) with the correct cell type specificity. Similarly, early spermatogenesis markers did appear to be differentially expressed in GSC/Early spermatogonia cell types, but with highly attenuated signals. For example, of all 1,326 significantly differentially expressed genes in GSC/Early spermatogonia, the most differentially expressed

classic marker gene was *His2Av*, which was ranked 375 when sorted by log(fold change), while the next closest marker was *aub*, ranked 600. This attenuation of signal appears to continue through later stages of spermatogenesis but reverses at the late spermatid stage. There, *p-cup* is ranked 18/639 and *soti* is ranked 24/639 respectively. Interestingly, cells demonstrating the “*ananassae*-specific” expression pattern showed significant overexpression of both early and late spermatogenesis markers: *His2Av* (ranked 245/366) and *p-cup* (ranked 73/366), respectively. The attenuation of cell-type information of previously utilized marker genes highlights how cell type-specific divergence (over evolutionary time) contributed to the masking of proper cell type assignment between species.

Among the 12 classic marker genes in our dataset, only *aly* appears in our 198-gene list, indicating that the remaining *D. melanogaster* marker genes’ ability to predict cell type likely occurs in a lineage-specific manner. Eight of the 12 marker genes did not appear in the 198-gene list as they showed high levels of species information, while the remaining four marker genes remained relatively similar across species. Of those, two genes (*aub*, *bam*) were excluded for having undefined F_i (cell type) values resulting from little to no variance in certain cell types. This low/no variance occurs due to having uniformly few to no counts within different cell types. Interestingly, the remaining two genes (*Hsp23*, *MtnA*) were not excluded for having evolutionarily diverged expression patterns, but they were excluded for having poor predictive ability for cell type. This appears to contradict the appearance of *MtnA* in an analysis of the most highly differentially expressed genes across cell types (regardless of species; see Dataset S2). This discrepancy arises from differences in statistical methodologies. *MtnA* expression is found only in a very small number of cells, primarily somatic cells. Differential expression analysis, which compares expression within one cell type vs. expression across all other cell types (“one-vs.-all”),

revealed elevated *MtnA* expression in somatic cells (all spp.) despite its low detection in many other cell types (all spp.). In contrast, ANOVA compares the between-cell type variance across all cell types, where *MtnA*’s limited detection results in a low F -statistic value due to minimal variance in most cell types. This highlights the value of performing cell type assignment on genes that show highly specific yet robust expression across cell types (Fig. 1G). This effect is particularly important in the context of scRNA-Seq technology, where the absence of detection may be as much the result of low read count as it is of biologically low expression. Focusing on quantitative differences within widely expressed genes like those in our 198-gene list, rather than a reliance on nondetection events of relatively lowly expressed genes, may lead to more consistent cell type classification in future comparative scRNA-Seq studies.

RNA In Situ Hybridization Chain Reaction Validates Cross-Species Cell Type Assignments.

While we were able to transfer a reference species’ cell type labels across species both in our simulated datasets and in our three-species dataset, it was unclear whether the cell type assignments reflected biological reality. To assess the accuracy of the cell type assignments, we performed in situ Hybridization Chain Reactions (HCR) RNA FISH (47–49) in whole-mount testis tissue from *D. melanogaster* and *D. ananassae*. Four genes were chosen: *Rbp4*, *B52*, *Pkd2*, and *soti* (Fig. 2) to fulfill multiple criteria: 1) highlighting predicted expression differences between species in common marker genes, 2) demonstrating high cell-type specificity to aid interpretation of subsequent in situ imaging, and 3) providing maximal information regarding our newly detected “*ananassae* spermatocyte” cluster. We note that during imaging, there was a large degree of crosstalk between the fluorophores representing *B52* and *soti*, so images were not amplified using both fluorophores in the same sample.




Fig. 2. Validation of expression pattern divergence and cell type classification using HCR RNA FISH. (A) scRNA-Seq UMAP shows predicted expression patterns in *D. melanogaster* and *D. ananassae* for four genes: *Rbp4*, *B52*, *Pkd2*, and *soti*. Key for cell type labels is found under *Rbp4*: Somatic, Germline stem cell (GSC)/Early spermatogonia, Late spermatogonia, *ananassae* spermatocyte, Early spermatocyte, Late spermatocyte, Early spermatid, Late spermatid. (B–E) RNA FISH images in *D. melanogaster* (B and C) and *D. ananassae* (D and E) using these same genes, with DAPI nuclear staining (B–E, Left column) and overlay of these channels over bright-field microscopy (B–E, Right column). *Rbp4* is a marker gene used in *D. melanogaster* to indicate germline stem cells/spermatogonia. Our scRNA-Seq cell type classification predicts that *Rbp4* has a broad expression pattern in *D. ananassae*. *B52* is a gene with a restricted expression pattern in *D. melanogaster* germline stem cells/spermatogonia. This expression pattern is predicted to be slightly broader in *D. ananassae*, extending into the new *ananassae* spermatocyte cluster. *Pkd2* is predicted to be primarily restricted to the newly identified *D. ananassae* spermatocyte cluster in both *D. melanogaster* and *D. ananassae*. *soti* is a marker gene used in *D. melanogaster* to indicate spermatid cell types, and our classification predicts that *soti* has both higher overall expression levels and lower cell type specificity in *D. melanogaster* than in *D. ananassae*.

Rbp4 was chosen as it was a previously employed cell-type marker for *D. melanogaster* GSC/Early spermatogonia cell types in prior publications (1). Our cross-species cell type classification predicted that this marker gene had highly differing expression patterns between *D. melanogaster* and *D. ananassae* (Fig. 2A). This prediction was validated, as *Rbp4* showed an unexpectedly wide expression pattern in *D. ananassae* beginning with early spermatogonia and culminating in expression overlap with late spermatids (*soti* expression and DAPI staining) (Fig. 2B–E). This is a surprising finding, as *Rbp4*, an RNA-binding protein, was found to be a regulator of mitosis. *Rbp4* interacts with *Fest* to regulate the translation of *Cyclin B* (50). Similarly, the genes *Lut* and *Syp* interact with *Fest*, regulating posttranscriptional control of *Cyclin B* (51). We then investigated the cell type-specific expression of the genes *Cyclin B*, *Fest*, and *Syp* in our dataset (SI Appendix, Fig. S6). *Lut* was excluded as it was not present as a one-to-one ortholog in our dataset. The expression pattern of *Cyclin B* agrees with previously reported expression patterns in *D. melanogaster* (50), while also adopting a similarly specific pattern in *D. yakuba*. Interestingly, the expression pattern of *Cyclin B* appears to be nonspecific in *D. ananassae*, showing a broad expression across all of spermatogenesis like *Rbp4*. While the *D. melanogaster* expression patterns of *Fest* and *Syp* were less specific than for *Rbp4* and *Cyclin B*, this low specificity decreased even further in *D. ananassae* (SI Appendix, Fig. S6). This result indicates that changes in cell type-biased patterns of genes potentially reflect underlying functional and molecular changes.

Finally, *soti* was chosen as it was previously employed as a cell-type marker for *D. melanogaster* late spermatid cell types (1, 2, 31, 35). Our cross-species cell type classification predicted that transcript levels in *D. melanogaster* would both be higher and less specific than in *D. ananassae*. This prediction was validated using our in situ imaging. For example, *D. melanogaster soti* expression appears to be broader than in *D. ananassae*, forming a broader, “smearier” expression pattern when compared to the tighter, more individualized puncta seen in *D. ananassae* (Fig. 2). Note that these figures show maximal intensity projections on z-stacks.

New Spermatocyte Cell Cluster Reveals a Unique Expression Pattern. A striking result of the previous sections is the identification of a *D. ananassae*-specific cluster that appears in our cross-species UMAP clustering. We thus use our scRNA-Seq and RNA FISH data to further investigate this newly identified cluster. As mentioned in the previous section, this “*ananassae*-specific” cluster appears in the UMAP of *D. ananassae* cells but does not seem well-represented in *D. melanogaster* and *D. yakuba* projections. While the appearance of this cluster seems correlated with the inferred disappearance of spermatid cell types in *D. ananassae*, the new cluster also appears in close proximity with other known spermatocyte cell types in our unified cross-species projections (Fig. 1D–F).

To resolve ambiguity and to further investigate the specificity of this new cluster, we identified genes that are differentially expressed (all spp.) in this cluster in comparison to all other clusters (*High Differential Expression across Cell Types in D. melanogaster Spermatogenesis Does Not Necessarily Indicate Differential Expression across Cell Types in Closely Related Species*). The 20 most highly differentially expressed genes in the *ananassae* spermatocyte cluster are reported in SI Appendix, Table S5.

While many of these differentially expressed genes are predicted to have expression in more than one tissue type, *Pkd2*, the 6th most differentially expressed gene in this cluster, was also found to have relatively specific expression. In particular, *Pkd2* demonstrated elevated cluster-specific expression in *D. ananassae*. While

specific, we note that *Pkd2* is still predicted to have residual expression from the GSC/Early spermatogenesis stage to the Early spermatid stages in both *D. melanogaster* as well as *D. ananassae* (Fig. 2A).

A detailed examination of the RNA FISH data for *Pkd2* reveals surprisingly broad expression throughout spermatogenesis (Fig. 2B–E), particularly in contrast to *B52* expression. *B52* was chosen because it is predicted to express in early spermatogonia as well as in our newly identified *ananassae*-specific cluster. As predicted, *B52* showed high specificity in germline stem cells and spermatogonia, with a residual expression detected in our *ananassae* spermatocyte cell cluster. These results confirm that cells with the *ananassae* spermatocyte expression pattern are a type of spermatocyte.

Expression of *Pkd2* in *D. melanogaster* and *D. ananassae* was found to overlap with GSC/Early spermatogonia as indicated by the expression pattern of *B52* (both species) and *Rbp4* (*D. melanogaster* only, “*Rbp4-mel*”). The expression of *Pkd2* subsequently decreases in concert with *B52* and *Rbp4-mel* as we continue along the germline differentiation process. Interestingly, *Pkd2* remains relatively highly expressed even in late-stage spermatocytes and onion-stage spermatids preceding spermatid elongation. This late-stage expression is most interestingly highlighted by the occasional presence of round cells that express *Pkd2* well into the spermatid stages of differentiation as represented by *soti* expression (Fig. 2B and E). This combination of scRNA-Seq and RNA FISH data suggests that the *ananassae* spermatocyte cell cluster has a biological component and is not the sole result of technical artifacts. However, we stress that this cluster does not likely represent a cell type that is functionally different from previously identified spermatocytes, but rather a shift in the expression pattern of a set of these cells. This is further discussed in SI Appendix, “*ananassae* Spermatocyte Cluster.”

Transcriptional Bursting in Spermatogenesis Is Controlled via Opposing Mechanisms during Meiotic Transition. We used our validated cell type assignments to study the evolution of transcriptional control by examining how the transcriptional bursting properties of different genes change during development. General transcriptional activity first increases and then dramatically decreases during the stage of spermatogenesis between the 16-spermatocyte stage and the second round of meiosis. These divisions subsequently result in 64 round spermatids in all three species (11). Importantly, the total pool of remaining transcripts is slowly degraded as they are translated in subsequent stages, supplemented only by a small number of genes that are transcribed postmeiotically. The increased resolution of scRNA-Seq analysis (26, 52) and MS2-MS2 coat protein system imaging (24, 25) has brought significant advances in understanding the fundamental molecular mechanisms underlying transcription, shedding light on the chemical kinetics that drive stochastic gene expression. However, global control of the pre- to postmeiotic transition during *Drosophila* spermatogenesis has remained understudied.

Aside from higher-order eukaryotic regulation via nucleosomes and histones, sequence-based transcriptional control is exerted by genomic segments that are broadly classified as promoter and enhancer sequences. Prior results have mechanistically demonstrated how this transcriptional control of promoters alters burst size via pol II availability, while control of enhancers mechanistically alters burst frequency via BRD4 and the Mediator complex (20, 26, 53–55). To investigate how transcriptional control of germline differentiation is controlled in *Drosophila*, we begin by calculating burst size and frequency for all annotated genes by cell type using “txburst,” using our three-species data (Fig. 3) (26). After applying default txburst quality control criteria, the number of genes that

were analyzed per cell type varied by cell type and species (*SI Appendix, Fig. S7 and Table S6*). Interestingly, we find that in the early, premeiotic stages of spermatogenesis (i.e., germline stem cells, early spermatogonia, late spermatogonia, and early spermatocytes), transcriptional bursting is primarily controlled via alterations in burst size. While burst frequency also increases during these stages, overall burst size appears to reach a maximum during the late spermatogonia stage before declining during spermatocyte stages (Fig. 3). Interestingly, in *D. melanogaster*, burst size drops sharply between the early and late spermatocyte stages, while in *D. yakuba* and *D. ananassae*, burst size decreases between the spermatogonia and spermatocyte stages, remaining relatively stable from early to late spermatocyte stages (Fig. 3 *A–F*). This genome-wide transition from burst size-dominant control (reflecting promoter-based regulation) to burst frequency-dominated control (reflecting enhancer-based regulation) suggests that phenotypic divergence during spermatogenesis in *Drosophila* may be evolving via global, genome-level alterations of transcriptional control via pol II initiation (k_{ini}) in alignment with prior observations (20, 53, 54, 56), rather than through alterations in a smaller subset of genes.

The control of bursting frequency appears to become the more dominant mechanism during postmeiotic stages of spermatogenesis (i.e., late spermatocyte, and spermatid stages, Fig. 3*B*). An increase in bursting frequency in *D. melanogaster* is observed in the late spermatocyte stage simultaneously with a correlated decrease in burst size. A similar change is observed between late spermatogonia and early spermatocyte in *D. yakuba*, where a decrease in burst size occurs simultaneously with an increase in bursting frequency (Fig. 3 *C and D*). In *D. ananassae*, an increase in burst frequency is observed in the early spermatocyte stage followed by an even larger increase in the late spermatocyte stage (Fig. 3 *E and F*). While the increase in burst frequency during the transition between late spermatogonia and early spermatocyte

stages is correlated with a decrease in burst size in the same stages, the second increase of burst frequency during the late spermatocyte stage does not show a correlated change in burst size. This increase in burst frequency is potentially an outlier, as not only did a very low number of annotated genes pass quality control measures (202 genes, *SI Appendix, Fig. S7C*), but also the variance in the observed distribution is also quite high. These observations could also reflect a deeper evolutionary divergence in transcriptomic control or cell type, resulting from a weaker cell type assignment as revealed by its sparse representation in UMAP and PCA projections (Fig. 1*E*). If the inferred distribution is closer in value to distributions found for the early spermatocytes and *D. ananassae* spermatocyte cell types, it would appear that *D. melanogaster* may have undergone a species-specific alteration where the size-to-frequency control transition has been delayed from the early spermatocyte stage to the late spermatocyte stage.

Overall, these results are consistent with a model of transcriptional control where an abundance of pol II is present (20, 26, 53, 54, 56) during earlier stages of spermatogenesis and becomes depleted during the meiotic transition. During the period of high overall pol II availability and activity, general levels of transcription remain high (20, 26, 53, 54, 56), and bursting kinetics are dominated by large burst size. Subsequent to this depletion, transcription then decreases significantly, shifting global transcriptional control into a more burst frequency-dominated mode of activity. It is likely that this modulation of burst frequency is achieved by enhancer-related changes in transcription factor stoichiometry (23, 25, 57).

Bursting Kinetics of X-Linked vs. Autosomal Genes Reveals That Partial Dosage Compensation Likely Occurs in Somatic and Germline Tissue. To help elucidate the timing of dosage compensation during spermatogenesis, we examined the transcriptional bursting kinetics for both X-linked ($N = 2,198$)




Fig. 3. Pre- and postmeiotic cell types in spermatogenesis are controlled by opposing mechanisms. The burst size and burst frequency distributions for (*A* and *B*) *D. melanogaster*, (*C* and *D*) *D. yakuba*, and (*E* and *F*) *D. ananassae* reveal that transcription in the earlier stages of spermatogenesis are primarily controlled via burst size, while transcription in later stages of spermatogenesis is primarily controlled via burst frequency. This transition between control mechanisms correlates with the transition between mitosis and meiosis.

and autosomal ($N = 11,749$) protein-coding genes. Interestingly, our dataset shows a highly consistent pattern where transcriptional burst size and frequency are significantly lower for X-linked genes than for autosomal genes across cell types and species (Mann–Whitney U test, Benjamini–Hochberg correction, $FDR < 0.05$) (Fig. 4). Significant differences between X-linked and autosomal bursting kinetics remain consistent throughout spermatogenesis. Perhaps most surprisingly, this difference is detected in *D. melanogaster* somatic tissue. While X-linked genes typically show lower activity than autosomal genes (even in statistically insignificant comparisons), the transcriptional bursting frequency for X-linked genes is higher than for autosomal genes in *D. yakuba* early spermatocytes. Additionally, while nearly every cell type in *D. melanogaster* and *D. yakuba* produces a significant difference between X- and autosomal-linked genes' transcriptional kinetics, the observed differences in *D. ananassae* are typically not significant after multiple hypothesis correction. This is likely because the number of genes that pass quality control in txburst is much lower for *D. ananassae* than for *D. melanogaster* and *D. yakuba*, resulting from the relatively poor reference genome quality and read depth for *D. ananassae* vs. *D. melanogaster* and *D. yakuba* (SI Appendix, Table S1).

Despite these differences, the ratio between X-linked and autosomal genes passing quality-control remains relatively constant across all stages and species (26) (Fig. 4 and SI Appendix, Fig. S7), reflecting both the number of genes that are expressed in each stage as well as read depth. The consistent ratio of X to autosomal genes expressed does not reveal a particular stage of spermatogenesis during which the entire X chromosome is entirely inactivated. After either the early spermatocyte stage in *D. melanogaster* or the

late spermatogonia stage in *D. yakuba* and *D. ananassae*, there is a drop-off in the number of genes that pass QC (SI Appendix, Fig. S7). Despite this change, the number of genes expressed on the X vs. autosomal chromosomes remains at a relatively constant ratio. As this ratio remains constant even when comparing somatic cell types to germline cell types (Fig. 4), it is possible that the process of X-downregulation occurs prior to the establishment of somatic vs. germline identity during the differentiation of testis tissue (58). This pattern could potentially be further tested by comparing the transcriptional bursting characteristics of X-linked to autosomal genes in earlier stages of testis organogenesis, e.g., in larval testis tissue.

Origination of New Transcripts Alters Regulation of Neighboring Genes by Increasing Burst Size during Meiotic Transition. The cultivator model of de novo gene origination hypothesizes that the most evolvable method of altering the expression of preexisting genes is the appearance of a new proximal promoter element (42). Unlike changes to existing transcription-factor-binding sites, as enhancer sequences typically act on longer range distances (≥ 1.5 kb), the cultivator model posits that the appearance of a promoter would act on relatively short ranges (~ 100 to 500 bp), likely through enhancer-mediated promoter interference (24) or supercoiling-mediated transcriptional coupling (59). One key prediction of this model would be that ncRNAs are found to be closer to their protein-coding neighbors than expected by chance—this is indeed the case (42). Another key prediction of this model is that the origination of a de novo transcript would alter the regulation of its immediate neighboring genes, but not more distant neighbors as might be expected by enhancers. To




Fig. 4. X-linked genes show lower bursting kinetics than autosomal genes prior to spermatogenesis. The burst size and burst frequency distributions for (A and B) *D. melanogaster*, (C and D) *D. yakuba*, and (E and F) *D. ananassae* reveal a conserved pattern of smaller average burst sizes and lower bursting frequencies for X-linked than autosomal genes across cell types (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). Note these are p-values before Benjamini–Hochberg correction, $FDR < 0.05$). No stage of spermatogenesis shows lower burst kinetics for X-linked genes than autosomal genes. The significant repression of X-linked genes in somatic cells suggests that decreased meiotic X chromosome expression occurs in a developmental stage preceding testis maturation.

test this prediction, we examined the transcriptional bursting characteristics of protein-coding genes that are neighboring newly activated de novo transcripts.

We begin by identifying four neighboring protein-coding genes for all 170 transcripts identified through bulk RNA-Seq analysis (Dataset S3), two upstream and two downstream (“L2,” “L1,” “R1,” and “R2” respectively, Fig. 5). Note that the L2 and L1 genes (or “5’-2” and “5’-1”) are defined as being in the 5’ direction of the de novo transcript, relative to its strand orientation—that is, upstream with respect to the transcript’s own direction. Similarly,

R2 and R1 genes (or “3’-1” and “3’-2”) are defined to be to the 3’ direction. Using these neighboring genes’ orthologous genes in *D. yakuba* and *D. ananassae*, we compared the burst size and frequency for each class of neighbors to genome-wide distributions. Unlike the 170 evolutionarily young de novo transcripts identified through our comparative analysis of bulk RNA-Seq data, the neighboring protein-coding genes are not expected to be biased in an evolutionary age.

Consistent with the cultivator model, the origination of these de novo transcripts is associated with a significant increase in




Fig. 5. Higher burst sizes in L1 neighboring genes of de novo transcripts support a cultivator model of de novo gene origination. (A) The initial step of the model suggests that a new promoter element near a preexisting gene may serve to regulate nearby genes in *cis*. We tested this by comparing the bursting kinetics of genes immediately neighboring new de novo transcripts to further neighboring genes. Note that in our study, de novo transcripts include those unique to *D. melanogaster* and those shared across the *melanogaster-simulans* or the *melanogaster-yakuba* clades. (B–D) L1 neighboring genes just upstream of de novo transcripts show significantly higher burst sizes than the distribution of all genes during the early and late spermatocyte stages, aligning with the meiosis onsite. These differences remain significant after Benjamini–Hochberg correction (FDR < 0.05). No other significant differences were observed in further neighboring genes or within spermatocytes preceding the de novo transcript origination (SI Appendix, Fig. S8).

transcriptional burst size in their 1st upstream neighboring genes, but not further neighboring genes. These differences are also found concurrently with the meiotic transition, suggesting that these genes may be functionally important in later stages of spermatogenesis. More specifically, significant differences are found either in the early spermatocyte (*D. melanogaster*, $P = 5.17 \times 10^{-4}$ and *D. yakuba*, $P = 4.56 \times 10^{-4}$, Mann–Whitney *U* test) and late spermatocyte (*D. yakuba*, $P = 8.25 \times 10^{-4}$) stages. These results remain significant after correction for multiple hypothesis testing using a Benjamini–Hochberg procedure (4 neighbors $\times$ 8 clusters $\times$ 3 species = 96 hypotheses, FDR < 0.05; Fig. 5 and *SI Appendix*, Figs. S8 and S9). Importantly, the cultivator model also predicts that there is no significant difference in transcriptional bursting kinetics prior to the origination of these de novo transcripts as these neighboring genes predate the appearance of the new de novo transcripts in the outgroup species, *D. ananassae*. Consistent with this hypothesis, no significant differences in burst size are detected in any neighboring genes. Furthermore, no significant differences are detected in burst frequency in any neighbors at any stage in any species. Given an increase in burst size but no alteration of burst frequency, these results suggest that a promoter-based effect, rather than an enhancer-based mechanism, drives the *cis*-regulatory effect of new promoter birth. We note that while it is possible that de novo transcripts appear next to older genes that already have large burst size kinetics, this does not explain why these neighboring genes show significantly elevated burst size after, but not before, de novo transcript origination.

To further validate this finding, we compared the genomic distribution of mean read count per cell (i.e. expression) with the same distributions calculated for L2, L1, R1, and R2 neighboring gene groups (*SI Appendix*, Fig. S10 and Table S7). While many significant differences in mean read count per cell (cell type–specific expression) were found, most of these significant differences persisted between in-group (*D. melanogaster* and *D. yakuba*) and outgroup (*D. ananassae*) comparisons or had opposing directions between species (e.g., significantly increased in *D. melanogaster* and decreased in *D. yakuba*). Only three neighboring group/cell type combinations showed significant differences within in-group species (*D. melanogaster* and *D. yakuba*) while also being directionally consistent: L1 neighbors in early spermatocytes (increase), R1 neighbors in early spermatocytes (decrease), and R1 neighbors in late spermatids (decrease).

In alignment with our previous finding of increased burst size, we find significantly elevated expression of L1 neighboring genes in early spermatocytes in *D. melanogaster* and *D. yakuba* but not *D. ananassae* (*SI Appendix*, Fig. S10). Interestingly, we also find significantly decreased expression of R1 neighboring genes in both early spermatocytes and early spermatids when compared to genomic expression. The combination of increased L1 neighboring gene expression as well as decreased R1 neighboring gene expression is particularly striking as it is consistent with supercoiling-mediated coupling (59). The appearance of a new promoter element in the tandem upstream/downstream conformations as shown in Fig. 5A would be expected to generate underwinding just upstream of and overwinding just downstream of the new promoter. This would result in both increased transcription in the upstream gene (the L1 neighbor), and decreased transcription of the downstream gene (the R1 neighbor).

Discussion

Feature Selection through ANOVA Allows for the Identification of Conserved Cell Type–Specific Genes. Our work demonstrates how the identification of the evolutionarily conserved 198-gene list allowed for cell type classifications from *D. melanogaster* to be extended to a panel of closely related *Drosophila* species. While it

was utilized in this study to overcome cell type–specific evolution, the methodology used here should be generalizable to a large range of problems where conditions and transcriptome expression vary in a correlated manner. For example, strong cell type–specific transcriptome-wide responses to external treatment (e.g., heat shock) could obscure cell type assignments. The application of an ANOVA-based methodology could be utilized to identify genes that differ strongly between cell types but remain robust to treatment.

Interestingly, the 198-gene list that we identified did not show a large overlap with more traditionally utilized cell type markers in *D. melanogaster* while still being significantly enriched for previously identified spermatogenesis-related function. One potential interpretation of this observation is that while the markers in the 198-gene list do have known cell type–specific function, it is possible that they do not perform as well as other, more classic markers within *D. melanogaster*. Alternatively, that the list of classic marker genes might reflect the historical discovery process regarding *D. melanogaster* spermatogenesis alone rather than underlying biological functions within the larger *melanogaster* species group. Regardless of the reason, our work has shown that the utilization of these marker genes in *D. melanogaster* does not necessarily imply that they are the most optimal markers when examining multiple species' genomes.

Biological Differences in Cell Type Classification. When we examined the *ananassae* spermatocyte cluster, we found biological evidence that there may be meaningful transcriptional differences in these cells compared to other cell types. For example, while these cells are poorly represented in *D. melanogaster* and *D. yakuba*, we found unusual patterns in our differential expression analysis in our full dataset. Recall that we found significant differential expression of *His2Av*, an early spermatogenesis marker in *D. melanogaster*, as well as *p-cup*, a late spermatogenesis marker in *D. melanogaster*. Consistent with this observation, we also note the early expression of *Pkd2* in GSC/Early spermatogonia, as well as the occasional presence of cells expressing *Pkd2* well into spermatid differentiation (Fig. 2 C and E). Notably, neither *His2Av* nor *p-cup* are found in our 198-gene list used for classification. Such transcriptional differences during the spermatocyte stage, combined with the detection of the “*ananassae*-specific spermatocyte” cluster, and unique *D. ananassae* reproductive biology during meiosis (60), suggests that the rapid evolution of the testis may be dominated by changes during this stage, reminiscent of earlier observations that de novo genes tend to express most highly in spermatocytes (35).

We also found that *Rpb4* has a surprisingly broad expression pattern in *D. ananassae*. This provokes the interesting question of whether this broad expression pattern is reflective of the ancestral state or evolved in an ancestor of *D. ananassae* subsequent to *D. melanogaster* speciation. It is likely that investigation into the upstream transcriptional differences that drive the alteration of *Rpb4* expression would lead to more general insight into the evolution of transcriptional programming in the testis. Similarly, investigation into the downstream effects of broad *Rpb4* expression would likely be fruitful. *Rpb4*, in conjunction with *fest* and *Syp*, has been shown to regulate *Cyclin B* during meiosis in *D. melanogaster* (50, 51). However, *Rpb4* and *Cyclin B* have much higher cell type specificity in *D. melanogaster* than *D. ananassae*, but in slightly different tissue types. How did these expression patterns evolve? Is the evolutionary appearance of this specificity coupled between the two genes? Alternatively, did this loss of specificity occur only in the *D. ananassae* lineage? These questions require further investigation in the future, particularly with the utilization of additional outgroup species.

We also note that our cross-species categorization detected an absence of spermatids in our scRNA-Seq data. However, our in situ

results showed highly specific expression of *soti*, while our bright-field microscopy revealed cells that can clearly be identified as late spermatids based on morphology alone. Interestingly, the gross morphology of these *D. ananassae* spermatids appeared to diverge from *D. melanogaster* as well, as they appeared to be “stringier” and “fuzzier” in addition to downstream morphological differences observed in the accessory gland, where tissue appeared to be larger and whiter. As the total amount of transcripts becomes increasingly depleted as spermatids fully mature, the transcriptional programming controlling these phenotypic differences likely occurs in earlier stages, e.g., “*ananassae* spermatocyte.” Consequently, the downstream biological effects of this altered transcriptional programming will likely be opaque to transcription-based techniques, highlighting the need for proteomic analysis or the application of traditional genetic tools beyond *D. melanogaster*.

Transcriptional Bursting Kinetics Offers Insight into Male Sex Chromosome Regulation. Many unanswered questions remain regarding the mechanisms that drive decreased X chromosome expression during spermatogenesis, leading to a large variety of proposed models. For example, differential polymerase elongation rates have been suggested as a mechanism underlying dosage compensation in males (61). Alternatively, a more canonical pathway involving the lncRNAs *roX1*, *roX2*, and the male-specific lethal complex (MSL) is possible (62, 63). One attractive aspect of the canonical MSL pathway hypothesis is that recent results have shown that the intrinsically disordered C-terminal region of *MSL2* and the *roX* lncRNAs induce X chromosome compartmentalization that functions similarly to phase separation (64). This has led to the suggestion that this X chromosome compartment could be enriched for pol II, resulting in increased initiation of transcription and an increase in transcriptional burst size (27). However, it has also been suggested that elements of the MSL complex, such as *maleless*, do not associate with the X chromosome during spermatogenesis (65).

Our results demonstrate a clear transition from a burst size-dominated mode of regulation to a burst frequency-dominated mode during the meiotic transition, suggesting that this transition may be driven by a global decrease in polymerase availability. However, in alignment with a prior analysis of previous results (31), our analysis of transcriptional bursting kinetics demonstrates that significantly decreased burst size and burst frequency for X-linked genes precedes not only the meiotic transition, but possibly also the establishment of germline stem cells. This decreased burst size is consistent with a study demonstrating decreased pol II binding on X chromosomes derived from both isolated spermatocytes and dissociated testis in comparison to wing imaginal discs (66). While these results are consistent with some manner of incomplete dosage compensation (31, 67), both decreased burst size and burst frequency suggest that a depletion of pol II localized

to the X chromosome alone is not a complete explanation for the observed X-linked decrease in expression. More specifically, a local depletion of pol II would decrease transcriptional burst size without altering burst frequency as observed in the developed embryo (27). Instead, the as-yet-unknown mechanism for decreased X chromosome expression should affect both the binding rate and the initiation rate of pol II to X chromosome-linked promoter sequences.

Another interesting result from our analyses is species-specific differences in kinetic bursting parameter values. The txburst software performs inference over raw read counts, and so it is likely that these differences are related to differences in read depth (*SI Appendix*, Fig. S10). Differences in transcript decay rates across species may also be a contributing factor (53). These differences are unlikely to reflect true biological differences in transcriptional burst kinetics, as these differences appear to be unbiased in regard to transcript identity (e.g., housekeeping genes). Importantly, we do not compare kinetic parameters across species, so such differences do not affect our conclusions; further experimental evidence, e.g., single-molecule studies, would be needed for proper cross-species comparisons.

Methods and Materials

Information on Methods and Materials used is available in *SI Appendix*. Briefly, we generated scRNA-Seq data from the testis of *D. melanogaster*, *D. yakuba*, and *D. ananassae*. ANOVA was used to transfer cell type classifications across species and applicability was validated by simulation and HCR RNA FISH (*SI Appendix*, Table S8). De novo transcripts were computationally identified, and bursting kinetics were inferred using maximum likelihood (26).

Data, Materials, and Software Availability. scRNA-Seq reads were uploaded to NCBI under the project number PRJNA995212 (68). Scripts for analyses may be found at https://github.com/LiZhaoLab/denovo_bursting_2025 (69). An R package for H_{50} implementation may be found at <https://github.com/ulee-sciscrpts/schSQ> (70).

ACKNOWLEDGMENTS. We thank Helen Duan and Connie Zhao from The Rockefeller University Genomics Resource Center for the help with scRNA-Seq library preparation, RRID:SCR_020986. Confocal microscopy was performed in the Rockefeller University's Bio-Imaging Resource Center, RRID:SCR_017791. We thank the Zhao lab members for helpful discussion and Sasha Mills for editing help. U.L. would like to thank Huang Suqi for helpful discussions regarding statistics. This work was supported by NIH MIRA R35GM133780, the Robertson Foundation, and an Allen Distinguished Investigator Award from Paul G. Allen Family Foundation to L.Z., and NSF postdoctoral fellowship 2410289 to U.L. The content of this study is solely the responsibility of the authors and does not necessarily represent the official views of the funders.

1. A. A. Raz *et al.*, Emergent dynamics of adult stem cell lineages from single nucleus and single cell RNA-Seq of *Drosophila* testes. *Elife* **12**, e82201 (2023).
2. E. Witt, C. B. Langer, N. Svetec, L. Zhao, Transcriptional and mutational signatures of the *Drosophila* ageing germline. *Nat. Ecol. Evol.* **7**, 440–449 (2023).
3. S. V. Nuzhdin, M. L. Wayne, K. L. Harmon, L. M. McIntyre, Common pattern of evolution of gene expression level and protein sequence in *Drosophila*. *Mol. Biol. Evol.* **21**, 1308–1317 (2004).
4. S. Khodursky, N. Svetec, S. M. Durkin, L. Zhao, The evolution of sex-biased gene expression in the *Drosophila* brain. *Genome Res.* **30**, 874–884 (2020).
5. Y. Hao *et al.*, Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573–3587.e29 (2021).
6. L. Haghverdi, A. T. L. Lun, M. D. Morgan, J. C. Marioni, Batch effects in single-cell RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat. Biotechnol.* **36**, 421–427 (2018).
7. I. Korsunsky *et al.*, Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289–1296 (2019).
8. A. J. Tarashansky *et al.*, Mapping single-cell atlases throughout Metazoa unravels cell type evolution. *Elife* **10**, e66747 (2021).
9. R. R. Snook, T. A. Markow, T. L. Karr, Functional nonequivalence of sperm in *Drosophila pseudoobscura*. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 11222–11226 (1994).
10. D. Joly, A. Korol, E. Nevo, Sperm size evolution in *Drosophila*: Inter- and intraspecific analysis. *Genetica* **120**, 233–244 (2004).
11. L. Schärer, J.-L. Da Lage, D. Joly, Evolution of testicular architecture in the *Drosophilidae*: A role for sperm length. *BMC Evol. Biol.* **8**, 143 (2008).
12. Y. M. Yamashita, M. T. Fuller, D. L. Jones, Signaling in stem cell niches: Lessons from the *Drosophila* germline. *J. Cell Sci.* **118**, 665–672 (2005).
13. C. Lim, L. Tarayrah, X. Chen, Transcriptional regulation during *Drosophila* spermatogenesis. *Spermatogenesis* **2**, 158–166 (2012).
14. Z. A. Alvi, T.-C. Chu, V. Schawaroch, A. V. Klaus, Protamine-like proteins in 12 sequenced species of *Drosophila*. *Protein Pept. Lett.* **20**, 17–35 (2013).
15. C.-H. Chang, I. Mejia Natividad, H. S. Malik, Expansion and loss of sperm nuclear basic protein genes in *Drosophila* correspond with genetic conflicts between sex chromosomes. *Elife* **12**, e85249 (2023).
16. H. H. McAdams, A. Arkin, Stochastic mechanisms in gene expression. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 814–819 (1997).
17. A. Raj, A. van Oudenaarden, Nature, nurture, or chance: Stochastic gene expression and its consequences. *Cell* **135**, 216–226 (2008).

18. I. Golding, J. Paulsson, S. M. Zawilski, E. C. Cox, Real-time kinetics of gene activity in individual bacteria. *Cell* **123**, 1025–1036 (2005).
19. L. Cai, N. Friedman, X. S. Xie, Stochastic protein expression in individual cells at the single molecule level. *Nature* **440**, 358–362 (2006).
20. J. P. Bothma *et al.*, Dynamic regulation of eve stripe 2 expression reveals transcriptional bursts in living *Drosophila* embryos. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 10598–10603 (2014).
21. A. Raj, C. S. Peskin, D. Tranchina, D. Y. Vargas, S. Tyagi, Stochastic mRNA synthesis in mammalian cells. *PLoS Biol.* **4**, e309 (2006).
22. D. M. Suter *et al.*, Mammalian genes are transcribed with widely different bursting kinetics. *Science* **332**, 472–474 (2011).
23. M. C. Walters *et al.*, Enhancers increase the probability but not the level of gene expression. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 7125–7129 (1995).
24. T. Fukaya, B. Lim, M. Levine, Enhancer control of transcriptional bursting. *Cell* **166**, 358–368 (2016).
25. B. Zoller, S. C. Little, T. Gregor, Diverse spatial expression patterns emerge from unified kinetics of transcriptional bursting. *Cell* **175**, 835–847. e25 (2018).
26. A. J. M. Larsson *et al.*, Genomic encoding of transcriptional burst kinetics. *Nature* **565**, 251–254 (2019).
27. L. Forbes Beadle, H. Zhou, M. Rattray, H. L. Ashe, Modulation of transcription burst amplitude underpins dosage compensation in the *Drosophila* embryo. *Cell Rep.* **42**, 112382 (2023).
28. M. D. Vbranovski, Meiotic sex chromosome inactivation in *Drosophila*. *J. Genomics* **2**, 104–117 (2014).
29. W. Hense, J. F. Baines, J. Parsch, X chromosome inactivation during *Drosophila* spermatogenesis. *PLoS Biol.* **5**, e273 (2007).
30. C. D. Meiklejohn, E. L. Landeen, J. M. Cook, S. B. Kingan, D. C. Presgraves, Sex chromosome-specific regulation in the *Drosophila* male germline but little evidence for chromosomal dosage compensation or meiotic inactivation. *PLoS Biol.* **9**, e1001126 (2011).
31. E. Witt, N. Svetec, S. Benjamin, L. Zhao, Transcription factors drive opposite relationships between gene age and tissue specificity in male and female *Drosophila* gonads. *Mol. Biol. Evol.* **38**, 2104–2115 (2021).
32. E. Witt, Z. Shao, C. Hu, H. M. Krause, L. Zhao, Single-cell RNA-sequencing reveals pre-meiotic X-chromosome dosage compensation in *Drosophila* testis. *PLoS Genet.* **17**, e1009728 (2021).
33. C. Kemkemer, W. Hense, J. Parsch, Fine-scale analysis of X chromosome inactivation in the male germ line of *Drosophila melanogaster*. *Mol. Biol. Evol.* **28**, 1561–1563 (2011).
34. M. Long, N. W. VanKuren, S. Chen, M. D. Vbranovski, New gene evolution: Little did we know. *Annu. Rev. Genet.* **47**, 307–333 (2013).
35. E. Witt, S. Benjamin, N. Svetec, L. Zhao, Testis single-cell RNA-seq reveals the dynamics of de novo gene transcription and germline mutational bias in *Drosophila*. *Elife* **8**, e47138 (2019).
36. M. T. Levine, C. D. Jones, A. D. Kern, H. A. Lindfors, D. J. Begun, Novel genes derived from noncoding DNA in *Drosophila melanogaster* are frequently X-linked and exhibit testis-biased expression. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 9935–9939 (2006).
37. S. Chen, Y. E. Zhang, M. Long, New genes in *Drosophila* quickly become essential. *Science* **330**, 1682–1685 (2010).
38. A. M. Gubala *et al.*, The Goddard and Saturn genes are essential for *Drosophila* male fertility and may have arisen de novo. *Mol. Biol. Evol.* **34**, 1066–1082 (2017).
39. L. Zhao, P. Saelao, C. D. Jones, D. J. Begun, Origin and spread of de novo genes in *Drosophila melanogaster* populations. *Science* **343**, 769–772 (2014).
40. J. M. Cridland, A. C. Majane, H. K. Sheehy, D. J. Begun, Polymorphism and divergence of novel gene expression patterns in *Drosophila melanogaster*. *Genetics* **216**, 79–93 (2020).
41. L. Zhao, N. Svetec, D. J. Begun, De novo genes. *Annu. Rev. Genet.* **58**, 211–232 (2024).
42. U. Lee, S. Mozeika, L. Zhao, A synergistic, cultivator model of de novo gene origination. *Genome Biol. and Evol.* **16**, evae103 (2024).
43. X. Qiu *et al.*, Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods* **14**, 979–982 (2017).
44. A. Suvorov *et al.*, Widespread introgression across a phylogeny of 155 *Drosophila* genomes. *Curr. Biol.* **32**, 111–123. e5 (2022).
45. H. Mi, A. Muruganujan, J. T. Casagrande, P. D. Thomas, Large-scale gene function analysis with the PANTHER classification system. *Nat. Protoc.* **8**, 1551–1566 (2013).
46. P. D. Thomas *et al.*, PANTHER: Making genome-scale phylogenetics accessible to all. *Protein Sci.* **31**, 8–22 (2022).
47. H. M. T. Choi *et al.*, Third-generation in situ hybridization chain reaction: Multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**, dev165753 (2018).
48. H. S. Bruce *et al.*, Hybridization Chain Reaction (HCR) in situ protocol (2021). <https://doi.org/10.17504/protocols.io.bunznv6>. Accessed 11 April 2025.
49. R. C. Thayer, E. S. Polston, J. Xu, D. J. Begun, Regional specialization, polyploidy, and seminal fluid transcripts in the *Drosophila* female reproductive tract. *Proc. Natl. Acad. Sci. U.S.A.* **121**, e2409850121 (2024).
50. C. C. Baker, B. S. Gim, M. T. Fuller, Cell type-specific translational repression of Cyclin B during meiosis in males. *Development* **142**, 3394 (2015).
51. C. C. Baker *et al.*, Cell-type-specific interacting proteins collaborate to regulate the timing of Cyclin B protein expression in male meiotic prophase. *Development* **150**, dev201709 (2023).
52. G. Hunt *et al.*, Tissue-specific RNA Polymerase II promoter-proximal pause release and burst kinetics in a *Drosophila* embryonic patterning network. *Genome Biol.* **25**, 2 (2024).
53. P. Trzaskoma *et al.*, 3D chromatin architecture, BRD4, and Mediator have distinct roles in regulating genome-wide transcriptional bursting and gene network. *Sci. Adv.* **10**, ead14893 (2024).
54. S. Gupta, J. J. Cai, Gene function revealed at the moment of stochastic gene silencing. *Commun. Biol.* **8**, 1–14 (2025).
55. M. Palacio, D. J. Taatjes, Real-time visualization of reconstituted transcription reveals RNA polymerase II activation mechanisms at single promoters. *bioRxiv [Preprint]* (2025). <https://www.biorxiv.org/content/10.1101/2025.01.06.631569v1> (Accessed 9 February 2025).
56. H. Ohishi *et al.*, Transcription-coupled changes in genomic region proximities during transcriptional bursting. *Sci. Adv.* **10**, eadn0020 (2024).
57. S. Alamos *et al.*, Minimal synthetic enhancers reveal control of the probability of transcriptional engagement and its timing by a morphogen gradient. *Cell Syst.* **14**, 220–236. e3 (2023).
58. S. Mahadevaraju *et al.*, Dynamic sex chromosome expression in *Drosophila* male germ cells. *Nat. Commun.* **12**, 892 (2021).
59. C. P. Johnstone, K. E. Galloway, Supercoiling-mediated feedback rapidly couples and tunes transcription. *Cell Rep* **41**, 111492 (2022).
60. B. N. Singh, *Drosophila ananassae*: A species characterized by spontaneous male recombination in appreciable frequency. *J. Genet.* **99**, 12 (2020).
61. E. Larschan *et al.*, X chromosome dosage compensation via enhanced transcriptional elongation in *Drosophila*. *Nature* **471**, 115–118 (2011).
62. A. Franke, B. S. Baker, The rox1 and rox2 RNAs are essential components of the compensasome, which mediates dosage compensation in *Drosophila*. *Mol. Cell* **4**, 117–122 (1999).
63. T. Conrad, A. Akhtar, Dosage compensation in *Drosophila melanogaster*: Epigenetic fine-tuning of chromosome-wide transcription. *Nat. Rev. Genet.* **13**, 123–134 (2012).
64. C. I. K. Valsecchi *et al.*, RNA nucleation by MSL2 induces selective X chromosome compartmentalization. *Nature* **589**, 137–142 (2021).
65. L. Rastelli, M. I. Kuroda, An analysis of *maleless* and histone H4 acetylation in *Drosophila melanogaster* spermatogenesis. *Mech. Dev.* **71**, 107–117 (1998).
66. J. T. Anderson, S. Henikoff, K. Ahmad, Chromosome-specific maturation of the epigenome in the *Drosophila* male germline. *Elife* **12**, RP89373 (2023).
67. K. H.-C. Wei, K. Chatla, D. Bachtrog, Single-cell RNA-seq of *Drosophila miranda* testis reveals the evolution and trajectory of germline sex chromosome regulation. *PLoS Biol.* **22**, e3002605 (2024).
68. U. Lee *et al.*, BioProject PRJNA995212 Single cell RNA sequencing of *Drosophila* testis tissue. NCBI Sequence Read Archive. <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA995212>. Deposited 15 July 2023.
69. U. Lee *et al.*, Code for Comparative single cell analysis of transcriptional bursting reveals the role of genome organization in de novo transcript origination. GitHub. https://github.com/LiZhaoLab/denovo_bursting_2025. Deposited 27 February 2025.
70. U. Lee, scHSQ. GitHub. <https://github.com/ulee-scscripts/scHSQ>. Deposited 4 February 2025.